

Quantitative Chemical Mass Transfer in Coastal Sediments during Early Diagenesis: Effects of Biological Transport, Mineralogy, and Fabric

Philippe Van Cappellen
Georgia Institute of Technology
221 Bobby Dodd Way
Atlanta, GA 30332 USA
phone (404) 894-3893 fax (404) 894-5638 email: pvc@eas.gatech.edu

Carla Koretsky
Georgia Institute of Technology
221 Bobby Dodd Way
Atlanta, GA 30332 USA
phone (404) 894-1579 fax (404) 894-5638
email: koretsky@eas.gatech.edu

Award # N00014-98-1-0203 (Grant #)
http://www.onr.navy.mil/sci_tech/ocean/onrpga.htm

LONG-TERM GOALS

Current multicomponent reactive transport models developed to study early diagenesis in nearshore environments are limited by their rudimentary mathematical descriptions of biologically-enhanced solute transport (bioirrigation) and biologically-enhanced particle transport (bioturbation). The long-term goal of this study is to develop models employing stochastic and inverse methods to quantify the spatial and temporal variation of bioirrigation and bioturbation in nearshore sediments and to incorporate information derived from these models into the multicomponent reaction transport model STEADYSED (Van Cappellen and Wang, 1996). This will greatly improve our ability to quantitatively describe early diagenesis in shallow-marine environments.

OBJECTIVES

The overall objective of this study was to quantify the effects of bioirrigation and biologically-induced mixing of particles on the early diagenesis of shallow-marine sediments. Specific objectives for FY99 included (1) the development of a stochastic model to describe bioirrigation intensity as a function of sediment depth in spatially and temporally heterogeneous environments using macrofaunal burrow densities and morphologies, (2) the development of an inverse nonlocal-transport model to describe bioirrigation intensity as a function of sediment depth based on vertical profiles of chemical species, and (3) the extraction of depth profiles of bioirrigation intensity for a field site using both the inverse and stochastic models.

APPROACH

An inverse, steady-state, 1-D reaction-transport model was developed with terms accounting for diffusion, advection, reaction and bioirrigation (e.g., Emerson et al., 1984; Boudreau, 1996),

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1999		2. REPORT TYPE		3. DATES COVERED 00-00-1999 to 00-00-1999	
4. TITLE AND SUBTITLE Quantitative Chemical Mass Transfer in Coastal Sediments during Early Diagenesis: Effects of Biological Transport, Mineralogy, and Fabric				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Georgia Institute of Technology, 221 Bobby Dodd Way, Atlanta, GA, 30332				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 7	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

$$\frac{\partial C}{\partial t} = \frac{\partial}{\partial z} \left(D \frac{\partial C}{\partial z} \right) - R - v \frac{\partial C}{\partial z} + \alpha (C_0 - C) = 0$$

where C is concentration of a given chemical species of interest, t is time, z is depth with respect to the sediment-water interface, D is the diffusion coefficient of the chemical species of interest, R is the depth-dependent rate of production or consumption of that species, v is the advective velocity, α is the depth-dependent bioirrigation coefficient and C_0 is the concentration of the species at the sediment-water interface. Depth-dependent bioirrigation coefficients were constrained with the model using sulfate concentration profiles measured in our laboratory and sulfate reduction rate profiles measured by J. Kostka (Skidaway Institute of Oceanography) for a series of sites at a saltmarsh on Sapelo Island, GA. Permeability measurements made by Y. Furukawa and D. Lavoie (Naval Research Laboratory) were used to constrain advective velocities in the sediments. A downhill simplex algorithm (Press et al., 1989) was used to minimize deviations between measured and calculated data and Monte Carlo simulations were used to avoid local minima. In addition, various vertical discretizations of the data were tested, so that the best regressed irrigation coefficient profile with the least number of fitting parameters could be extracted from the measured data.

A stochastic model of burrow distributions was developed to function as a link between ecological data and bioirrigation coefficients. This approach allows the extreme spatial and temporal heterogeneity of nearshore depositional environments to be considered explicitly in constraining solute transport via bioirrigation. The basic approach of the stochastic modeling is to create probabilistic descriptions of burrow densities and morphologies and to use those descriptions to simulate burrow networks under different ecological scenarios (i.e., for sediments with only one type of burrow or for sediments with a mixture of burrow types). The burrow networks are then used to calculate the number of burrows encountered as a function of depth (a proxy for the surface area of exchange) which is directly related to the bioirrigation coefficients.

WORK COMPLETED

An inverse model to minimize deviations between measured and calculated data was developed and used to extract seasonal irrigation coefficient profiles at three sites in a saltmarsh at Sapelo Island, GA.

A stochastic model was developed to relate the distribution and morphology of burrows to bioirrigation intensity (Koretsky et al., 1999). Burrow networks were simulated using probability functions derived from literature data (Basan and Frey, 1977; Teal, 1958) and from X-ray data collected by Y. Furukawa and D. Lavoie (NRL) for saltmarsh sites at Sapelo Island, GA. Simulated networks were used to calculate the depth-dependent surface area of exchange between burrow walls and the porous sediment matrix (a proxy for the bioirrigation coefficient).

RESULTS

The inverse model has been used to regress irrigation coefficient profiles for three sites (a creek bank, the adjacent levee and a ponded marsh site) in a saltmarsh at Sapelo Island, GA using sulfate concentration and reduction rate profiles measured in summer, spring and fall. Irrigation was always

found to be the *dominant solute transport process* at all three sites. Total calculated irrigation fluxes of dissolved sulfate are shown in Figure 1.




Figure 1. Integrated irrigation fluxes of sulfate for creek bank, levee and ponded marsh sites at Sapelo Island, GA for spring (solid), summer (hatched) and fall (stippled).

It can be seen in Figure 1 that irrigation varies strongly as a function of both site and season, with the greatest integrated irrigation flux occurring in summer at the levee site. Irrigation coefficient profiles showed that irrigation always decreases rapidly with depth at all sites.

Burrow networks were generated for these same three saltmarsh sites using the stochastic model. The resulting number of burrows intersected as a function of depth (a proxy for the bioirrigation intensity) and the associated standard deviation at the ponded marsh site are shown in Figures 2 and 3. These calculations suggest that the irrigation is most intense in the top ten to twenty centimeters of sediment, and that irrigation is deepest at the creek bank and levee sites and shallowest at the ponded marsh site. The model also suggests that irrigation is least intense at the ponded marsh site.




Figure 2. Number of burrows encountered as a function of depth for ponded marsh (squares), creek bank (diamonds) and levee (circles) sites calculated from stochastic model.




Figure 3. Number of burrows encountered as a function of depth for ponded marsh; error bars represent one standard deviation (based on 10,000 simulations).

IMPACT/APPLICATIONS

The inverse model developed in this study represents a significant improvement over previous nonlocal transport models (e.g., Emerson et al., 1984; Boudreau, 1984) because unbiased bioirrigation coefficient profiles may be extracted from chemical data, with explicit account taken of uncertainties in the measured data, and without imposing an a priori choice of the functional dependence of the

irrigation exchange coefficient with depth. In addition, the model may be used to extract production/consumption rate profiles from measured concentration data for chemical species other than sulfate, once the irrigation coefficient profiles have been derived.

Calculated surface areas of exchange from the stochastic model were found to be qualitatively consistent with irrigation profiles obtained independently using the inverse model. However, differences in profiles obtained from the two models suggest that the treatment of burrow water chemistry and fluid dynamics (flushing intensity and frequency) must be improved if the two models are to be linked. For example, the intensity of irrigation at the creek bank site predicted from the inverse model is considerably less than predicted by the burrow network simulations. This may be due to decreased flushing frequency at this site, associated with lower sulfide pore water concentrations, a result which could not be predicted by burrow network size alone.

TRANSITIONS

The inverse and stochastic models developed in this study will be used by Y. Furukawa (NRL) to constrain bioirrigation at field sites (Bay St. Louis; Horn Island, MS) and in experimental mesocosms (in collaboration with S. Bentley, Louisiana State University).

RELATED PROJECTS

Seasonal pore water profiles of dissolved species (e.g., Fe(II)/Fe(III), SO₄(-II), H₂S, and Mn) have been measured along a transect in a saltmarsh at Sapelo Island, GA (Koretsky et al., in prep). The seasonal oscillation of microbial community structure at these same sites was studied in collaboration with Dr. T. DiChristina (Georgia Institute of Technology). The bioirrigation data obtained using the inverse and stochastic models in this study will aid in interpreting the observed temporal and spatial relationships between pore water and sediment geochemical conditions and microbial community structure in the saltmarsh. In addition, a global relationship has been found between irrigation intensities and rates of organic carbon degradation that may be useful in constraining irrigation intensities in modern and ancient ocean sediments.

REFERENCES

- Basan P.B. and Frey R.W. (1977) Actual paleontology and neoichnology of salt marshes near Sapelo Island, Georgia. In Trace Fossils 2 (ed. T.P. Crimes and J.C. Harper), pp. 41-70, Steel House Press.
- Boudreau B. (1996) Diagenetic Models and Their Implementation. New York, Springer-Verlag.
- Emerson S., Jahnke R., and Heggie D. (1984) Sediment-water exchange in shallow water estuarine sediments. Jour. Mar. Res. 42, 709-730.
- Koretsky C.M., Van Cappellen P. and Kostka J. (1998) Seasonal dependence of porewater irrigation in a saltmarsh at Sapelo Island, GA: Implications for sediment biogeochemistry. Geological Society of America, Abstracts with Programs.
- Koretsky C.M., Van Cappellen P. and Meile C. (1999) A stochastic approach to bioirrigation modeling in sedimentary environments. Fifth International Symposium on the Geochemistry of the Earth's Surface. August 12-20, Reykjavik, Iceland.

- Meile C., Koretsky C., Van Cappellen P. and Kostka J. (1999) Spatial and temporal variation in bioirrigation: An inverse approach. Fifth International Symposium on the Geochemistry of the Earth's Surface. August 12-20, Reykjavik, Iceland.
- Press W.H., Flannery B.P., Teukolsky S.A. and Vetterling W.T. (1989) Numerical Recipes: The art of scientific computing (FORTRAN version). Cambridge University Press, New York, 702 pp.
- Teal J.M. (1958) Distribution of fiddler crabs in Georgia salt marshes. *Ecol.* 39(2), 185-193.
- Van Cappellen P. and Wang Y. (1996) Cycling of iron and manganese in surface sediments: a general theory for the coupled transport and reaction of carbon, oxygen, nitrogen, sulfur, iron, and manganese. *American Journal of Science* 296, 197-243.

PUBLICATIONS

- Koretsky C.M., Van Cappellen P. and Kostka J. (1998) Seasonal dependence of pore water irrigation in a salt marsh at Sapelo Island, GA: Implications for sediment biogeochemistry. *Geological Society of America, Abstracts with Programs*, 30, 374.
- Koretsky C.M. and Van Cappellen P. (1999) Spatial and temporal trends in salt marsh sediment geochemistry. 6th Symposium on Biogeochemistry of Wetlands. July 11-14, 1999. Ft. Lauderdale, Florida.
- Koretsky C.M. and Van Cappellen P. (1999) Quantitative Modeling of Bioirrigation: A Stochastic Approach. Chemical Oceanography Gordon Conference, Kimball Union Academy, August 8-13, 1999.
- Koretsky C.M., Van Cappellen P. and Meile C. (1999) A stochastic approach to bioirrigation modeling in sedimentary environments. 5th International Symposium on the Geochemistry of the Earth's Surface. August 16-20, 1999, Reykjavik, Iceland.
- Kostka J., Roychoudhury A., Koretsky C. and Van Cappellen P. (1999) Microbial respiration rates in saltmarsh sediments. 6th Symp. on the Biogeochemistry of Wetlands, July 11-14, 1999, Ft. Lauderdale, Florida.
- Lowe, K. L., DiChristina T. J., Roychoudhury A., and Van Cappellen, P. (1999) Microbial community structure and geochemistry of Sapelo Island salt marsh sediments. *Geomicrobiology Journal* (in press).
- Meile C., Koretsky C. and Van Cappellen P. (1999) Spatial and temporal variations in bioirrigation: An inverse approach. 5th International Symposium on the Geochemistry of the Earth's Surface. August 16-20, 1999, Reykjavik, Iceland.
- Meile, C. (1999) An inverse model for reactive transport in biogeochemical systems: Application to biologically-enhanced pore water transport (irrigation) in aquatic sediments. Georgia Institute of Technology (M.S. degree expected in Dec. 1999).
- Roychoudhury A.N. (1999) Biogeochemical dynamics in aquatic sediments: novel laboratory and field-based approaches. Georgia Institute of Technology (Ph.D. awarded spring, 1999).
- Van Cappellen P. (1999) Reactive transport in sediments and groundwater systems: What can we borrow from hydrologists (invited). Chemical Oceanography Gordon Conference, Kimball Union Academy, August 8-13.
- Van Cappellen P. (1999) Reactive transport modeling of complex biogeochemical systems (invited). 5th Society for Industrial and Applied Mathematics. Conference on Mathematical and Computational Issues in the Geosciences, San Antonio, March 24-27.

Van Cappellen P. and Steefel C. (1998) Advances in reactive transport modeling of field-scale environmental systems (invited). American Geophysical Union Fall Meeting, San Francisco, December 6-10.

Viollier, E., Inglett, P.W., Hunter, K., Roychoudhury, A.N., and Van Cappellen, P. (1999) The ferrozine method revisited: Fe(II)/Fe(III) determination in natural waters. Applied Geochemistry (in press).

PATENTS

none